

nextnano.QCL - Software documentation

General remarks

nextnano.QCL is a console application that is run from within the [nextnanomat](#) software (GUI). Alternatively, it can be executed from the command line. The input file specifies the device that shall be simulated.

nextnanomat

[nextnanomat](#) is a convenient graphical user interface for nextnano.QCL. It can be downloaded from [here](#). It can visualize 1D, 2D and 3D simulations results.

Input file

The input file specifies all properties of the device, such as geometry, material composition, temperature, grid,... Furthermore, it sets all parameters that are needed to define the program flow of nextnano.QCL. The keywords that can be used for this purpose are defined in the [syntax of the input file](#).

Output

nextnano.QCL exports its results to a directory. The output files are documented [here](#).

Examples

The nextnano.QCL installation provides some [example tutorials](#) that can be run with nextnanomat, to get familiar with the program.

Database

All material properties that are needed for simulation are described [here](#).

Download

The software can be obtained from [here](#).


If you have further questions, see the [FAQ](#) or contact support@nextnano.com.

Working principle

The code uses **field-periodic boundary condition**. In this way the simulation accounts for an infinite periodic structure, with a periodic electric field. Coherent transport between periods is accounted on a length set by `<Coherence_length_in_Periods>`.

The results of the calculation should not depend on which material layer the sequence starts, i.e. a cyclic permutation in the material layer sequence should not change the simulation results.


Program flow



In the beginning of the calculation, the single-band effective mass Schrödinger equation is solved in real space. The calculated energy levels and wave functions are then used as input to the NEGF algorithm. The wave functions are termed **modes** and the NEGF algorithm is written in terms of **mode space** and not **real space** to make it computationally more efficient. The number of QCL periods that are input to this Schrödinger equation are specified in `<Number_of_lateral_periods_for_band_structure>`. In general, the core of the NEGF algorithm should be rather independent of this number, e.g.

```
<Number_of_lateral_periods_for_band_structure> 4  
</Number_of_lateral_periods_for_band_structure>
```

should lead to very similar results compared to a value of 5 but the numerical values might differ slightly.



From:
<https://nextnano-docu.northeurope.cloudapp.azure.com/dokuwiki/> - **nextnano.NEGF - Software for Quantum Transport**

Permanent link:
https://nextnano-docu.northeurope.cloudapp.azure.com/dokuwiki/doku.php?id=qcl:software_documentation&rev=1489594147

Last update: **2017/03/15 16:09**